

ISSN 0840-8440

PROCEEDINGS

TECHNOLOGY TRANSFER CONFERENCE 1988

November 28 and 29, 1988

Royal York Hotel

Toronto, Ontario

SESSION A

AIR QUALITY RESEARCH

Sponsored by

Research and Technology Branch

Environment Ontario

Ontario, Canada

AASG.

Copyright Provisions and Restrictions on Copying:

This Ontario Ministry of the Environment work is protected by Crown copyright (unless otherwise indicated), which is held by the Queen's Printer for Ontario. It may be reproduced for non-commercial purposes if credit is given and Crown copyright is acknowledged.

It may not be reproduced, in all or in part, for any commercial purpose except under a licence from the Queen's Printer for Ontario.

For information on reproducing Government of Ontario works, please contact ServiceOntario Publications at copyright@ontario.ca

AP5

POLLUTANT TRANSPORT AND ACID RAIN FORMATION IN A FIELD OF CONVECTIVE CLOUDS: RESULTS FROM A MESOSCALE MODEL SIMULATION

Michał Niewiadomski
Department of Physics, University of Toronto
Toronto, Ontario, M5S 1A7

1. INTRODUCTION

Transport, scavenging and chemical transformations of sulphur and nitrogen pollutants in a field of convective clouds are studied in this paper by means of a three-dimensional mesoscale model, with horizontal resolution of 10 km and parameterized subgrid effects. Attention is focused on the average effects of cumulus convection, as well as the relative importance of different microphysical and chemical processes.

The paper is an outcome of the author's stay at the GKSS Forschungszentrum Geesthacht GmbH in Geesthacht, West Germany and the collaboration with its staff on the development of a mesoscale model of transport and transformations of atmospheric pollutants.

The University of Toronto and Ontario Hydro cloud chemistry model (Iribarne and Mele, 1988) has been coupled for this study with the developed at GKSS cloud microphysics module (Levkov et al., 1987) and a parameterization scheme for the subgrid effects of cumulus convection (Levkov et al., 1986a). These models are driven by a relatively simple dynamical model, which will be replaced in future with a sophisticated mesoscale model GESIMA being developed at GKSS. The model is briefly described below. More detailed description is given by Niewiadomski et al. (1989). Detailed formulas can be found in appropriate reports referenced in section 2.

2. THE MODEL

2.1 Dynamics

The dynamical module is based on the model of Beniston (1984). Detailed formulas can be found in papers of Levkov et al. (1986a,b). The model solves the hydrostatically approximated prognostic equations for momentum, potential temperature, specific humidity, hydrometeor mixing ratios, and chemical variables. The turbulent transport is parameterized through first-order closure; surface fluxes are computed according to Businger et al. (1971). Source terms due to microphysical and chemical processes as well as due to the subgrid effects of cumulus clouds are provided by the described below microphysical, chemical and cloud modules.

2.2 Cloud microphysics

The model follows the 'bulk microphysics' approach. The liquid water is divided into rain and cloud water fractions. Cloud ice and snow are combined in one variable, which will be referred to as 'snow'. The following microphysical processes are included: condensation, evaporation and freezing of cloud water, autoconversion of cloud water to rain, sublimational growth of snow, accretion of cloud droplets by rain and snow particles, freezing and evaporation of rain and melting of snow. Transfers of pollutants between cloud water, rain and snow due to microphysical processes are computed in the chemical module. Detailed formulas can be found in Levkov et al. (1987). Various aspects of this model are also discussed by Levkov et al. (1988).

2.3 Chemistry

The chemical module, computing the source terms for chemical variables due to chemical and microphysical processes is a slightly modified version of the model of Iribarne and Mele (1988). Its basic ideas are



(8018)
TD/5/T43

The chemical, microphysical and cumulus parameterization modules coupled for this study proved to be an effective tool for studying the mesoscale effects of cumulus clouds. After further improvements the modules will be incorporated in a more sophisticated dynamical model.

H_2O_2 was the main oxidant in cases studied, with sulphate production strongly dependent on the amount of H_2O_2 available. Sulphate production was depended also on the amount of liquid water in the domain. NO_x oxidation was of secondary importance in this study.

The average vertical profiles of SO_2 in air and cloud water are given Fig.3 for run 1 in 30 min intervals. The upward transport of SO_2 is clearly seen. The profiles resemble those obtained by Niewiadomski (1966) for a passive pollutant with a resolution of 1 km.

TABLE 2.

The sulphate budget, kg equivalents.

		run 1	run 2	run 3
Produced by SO_2 oxidation:				
in cloud water,	by H_2O_2	1510	5100	147
in cloud water,	by O_3	50	20	117
in rain,	by H_2O_2	7.9	0.9	2.6
in rain,	by O_3	0.2	0.5	0.9
cloud to rain transfer		598	851	237
cloud to snow transfer		23	0	8.5
below cloud scavenging		144	44	129
wet deposition		578	220	280

6. ACKNOWLEDGEMENT

The valuable discussions with Dr. Lubomir Levkov and his help in adapting for this study the dynamical and microphysical modules, as well as the help of Mr. Manuel Resch in preparing the plots for this paper are greatly appreciated.

7. REFERENCES

- Augstein, E. and W. Wendel, 1981: Modelling of the time-dependent atmospheric tradewind boundary layer. *Contr. Atmos. Phys.*, 53, 509-538.
- Beniston, M., 1984: A numerical study of atmospheric mesoscale cellular convection. *Dyn. Atmos. Ocean*, 8, 223-242.
- Businger, J.A., J.C. Wyngaard, Y. Izumi, and E.F. Bradley, 1971: Flux-profile relationships in the atmospheric surface layer. *J. Atmos. Sci.*, 28, 181-189.
- Iribarne, J.V. and H.-R. Cho, 1988: Models of cloud chemistry. *Tellus*, in press.
- Iribarne, J.V. and O.T. Melo, 1988: The role of clouds in the long-range transport of atmospheric pollutants and acid rain formation in Canada. Vol 3: Cloud chemistry. Canadian Electrical Association,

CEA No. 403 G 348.

Levkov, L., D. Eppel and H. Grassl, 1986a: On the simulation of convective structures. GKSS/86/E/35.

Levkov, L., D. Eppel and H. Grassl, 1986b: Simulation of convection over the ocean in the region of the intertropical convergence zone. GKSS 86/E/55.

Levkov, L., D. Eppel and H. Grassl, 1987: Parameterization of phase changes of water in a mesoscale model. GKSS 87/E/44.

Niewiadomski, M., 1986: A passive pollutant in a three-dimensional field of convective clouds: numerical simulations. *Atmospheric Environment*, 20, 139-145.

Niewiadomski, M., 1989: Sulphur dioxide and sulphate in a three-dimensional field of convective clouds: numerical simulations. *Atmospheric Environment*, in press.

Niewiadomski, M., L. Levkov, D. Jacob, and D. Eppel, 1989: Three-dimensional numerical simulations of pollutant transport and acid rain formation in a mesoscale field of convective clouds. To appear in preprints of the Symposium on the Role of Clouds in Atmospheric Chemistry and Global Climate, Anaheim, California, Jan. 23 - Feb. 3, 1989.




Fig. 3